

No stability switching at saddle-node bifurcations of solitary waves in generalized nonlinear Schrödinger equations

Jianke Yang

Department of Mathematics and Statistics, University of Vermont, Burlington, Vermont 05401, USA

(Received 28 November 2011; published 30 March 2012)

Saddle-node bifurcations arise frequently in solitary waves of diverse physical systems. Previously it was believed that solitary waves always undergo stability switching at saddle-node bifurcations, just as in finite-dimensional dynamical systems. Here we show that this is not true. For a large class of generalized nonlinear Schrödinger equations with real or complex potentials, we prove that stability of solitary waves does not switch at saddle-node bifurcations. This analytical result is confirmed by numerical examples where both soliton branches are stable at saddle-node bifurcations.

DOI: [10.1103/PhysRevE.85.037602](https://doi.org/10.1103/PhysRevE.85.037602)

PACS number(s): 05.45.Yv, 42.65.Tg

Saddle-node bifurcation is a well known phenomenon in finite-dimensional dynamical systems [1]. In this bifurcation, there are two fixed-point branches on one side of the bifurcation point and no fixed points on the other side, and the stability of these two fixed-point branches switches at the bifurcation point (one branch stable and the other branch unstable). In nonlinear partial differential equations (which can be viewed as infinite-dimensional dynamical systems), this bifurcation exists as well (it is also called fold bifurcation in the literature). For instance, solitary waves in nonlinear physical systems often exhibit this type of bifurcation. Examples include the Boussinesq equations and the fifth-order Korteweg–de Vries equation in water waves [2–4], the Swift-Hohenberg equation in pattern formation [5], the nonlinear Schrödinger (NLS) equations with localized or periodic potentials in nonlinear optics and Bose-Einstein condensates [6–9], and many others. Motivated by stability switching of saddle-node bifurcations in finite-dimensional dynamical systems, it is widely believed that in nonlinear partial differential equations, stability of solitary waves also always switches at saddle-node bifurcations (see [5–8] for examples). This belief is very pervasive since no counterexample has been reported yet.

In this paper, we show that this belief of universal stability switching at saddle-node bifurcations in nonlinear partial differential equations is incorrect. Specifically, we show that in a wide class of generalized NLS equations with real or complex potentials, stability of solitary waves actually does not switch at saddle-node bifurcations. This fact is proved analytically by using general conditions of saddle-node bifurcations and eigenvalue-bifurcation analysis. It is also verified numerically by several examples, where both branches of solitary waves are stable at saddle-node bifurcations.

We consider general nonlinear Schrödinger-type equations with arbitrary forms of nonlinearity and external potentials in multidimensions,

$$iU_t + \nabla^2 U + F(|U|^2, \mathbf{x})U = 0, \quad (1)$$

where ∇^2 is the Laplacian in the N -dimensional space $\mathbf{x} = (x_1, x_2, \dots, x_N)$, and $F(\cdot, \cdot)$ is a general function which contains nonlinearity as well as external potentials. These equations include the Gross-Pitaevskii equation in Bose-Einstein condensates and nonlinear light-transmission equations in localized or periodic potentials as special cases [10–12]. Below, we will

first focus on the case where the function F is real valued, which applies when the system (1) is conservative. Extension to the nonconservative case of complex functions of F will be considered afterward.

When the function F is real, Eq. (1) admits stationary solitary waves of the form

$$U(\mathbf{x}, t) = e^{i\mu t} u(\mathbf{x}), \quad (2)$$

where $u(\mathbf{x})$ is a localized real function satisfying

$$\nabla^2 u - \mu u + F(u^2, \mathbf{x})u = 0, \quad (3)$$

and μ is a real propagation constant which is a free parameter. Under certain conditions, these solitary waves undergo saddle-node bifurcations at special values of μ [6–9]. A signature of these bifurcations is that on one side of the bifurcation point μ_0 , there are no solitary wave solutions, but on the other side of μ_0 , there are two distinct solitary-wave branches which merge with each other at $\mu = \mu_0$. To derive conditions for these bifurcations, we introduce the linearization operator of Eq. (3),

$$L_1 = \nabla^2 - \mu + \partial_u [F(u^2, \mathbf{x})u]. \quad (4)$$

We also introduce the standard inner product of functions $\langle f, g \rangle = \int_{-\infty}^{\infty} f^*(\mathbf{x}) g(\mathbf{x}) d\mathbf{x}$, where the superscript “*” represents complex conjugation. Our analysis starts with the basic observation that, if a bifurcation occurs at $\mu = \mu_0$, by denoting the corresponding solitary wave and the linearization operator as

$$u_0(\mathbf{x}) = u(\mathbf{x}; \mu_0), \quad L_{10} = L_1|_{\mu=\mu_0, u=u_0}, \quad (5)$$

then the linear operator L_{10} should have a discrete zero eigenvalue. This is a necessary condition for all types of bifurcations. To derive sufficient conditions for saddle-node bifurcations, let us assume that this zero eigenvalue of L_{10} is simple, which is the case for generic bifurcations in one spatial dimension as well as for many bifurcations in higher spatial dimensions. Under this assumption, we denote this unique discrete (localized) real eigenfunction of L_{10} at the zero eigenvalue as $\psi(\mathbf{x})$, i.e.,

$$L_{10}\psi = 0. \quad (6)$$

We also denote

$$G(u; \mathbf{x}) = F(u^2; \mathbf{x})u, \quad G_2(\mathbf{x}) = \partial_u^2 G|_{u=u_0}. \quad (7)$$

Then the sufficient condition for saddle-node bifurcations of solitary waves is given by the following theorem.

Theorem 1. Under the above assumption and notations, if $\langle u_0, \psi \rangle \neq 0$ and $\langle G_2, \psi^3 \rangle \neq 0$, then a saddle-node bifurcation of solitary waves occurs at $\mu = \mu_0$ in Eq. (3).

Proof. Solitary waves which exist near $\mu = \mu_0$ admit the following perturbation series expansions:

$$u(\mathbf{x}; \mu) = \sum_{k=0}^{\infty} (\mu - \mu_0)^{k/2} u_k(\mathbf{x}). \quad (8)$$

Inserting this expansion into Eq. (3), we get the following equations for u_k at order $(\mu - \mu_0)^{k/2}$, $k = 0, 1, 2, \dots$:

$$\nabla^2 u_0 - \mu_0 u_0 + F(u_0^2, \mathbf{x}) u_0 = 0, \quad (9)$$

$$L_{10} u_1 = 0, \quad (10)$$

$$L_{10} u_2 = u_0 - G_2 u_1^2 / 2!, \quad (11)$$

and so on. Equation (9) for u_0 is satisfied automatically since u_0 is a solitary wave at $\mu = \mu_0$. The u_1 solution to Eq. (10) is found from Eq. (6) as

$$u_1 = b_1 \psi, \quad (12)$$

where b_1 is a constant. The u_2 function satisfies the linear inhomogeneous equation (11). Due to the Fredholm alternative theorem and the fact that L_{10} is self-adjoint, Eq. (11) admits a localized solution for u_2 if and only if the homogeneous solution ψ is orthogonal to the inhomogeneous term, i.e.,

$$\langle \psi, u_0 - G_2 u_1^2 / 2 \rangle = 0. \quad (13)$$

Inserting the solution (12) into this orthogonality condition and recalling the conditions in Theorem 1, we find that

$$b_1 = b_1^{\pm} \equiv \pm \sqrt{\frac{2\langle u_0, \psi \rangle}{\langle G_2, \psi^3 \rangle}}. \quad (14)$$

Thus we get two b_1 values b_1^{\pm} which are opposite each other. Inserting the corresponding u_1 solutions (12) into Eq. (8), we then get two perturbation-series solutions of solitary waves $u(\mathbf{x}; \mu)$ as

$$u^{\pm}(\mathbf{x}; \mu) = u_0(\mathbf{x}) + b_1^{\pm} (\mu - \mu_0)^{1/2} \psi(\mathbf{x}) + O(\mu - \mu_0). \quad (15)$$

If $\langle u_0, \psi \rangle$ and $\langle G_2, \psi^3 \rangle$ have the same sign, then b_1^{\pm} are real. Recalling that $u_0(\mathbf{x})$ and $\psi(\mathbf{x})$ are both real as well, we see that these perturbation-series solutions (15) give two real-valued (legitimate) solitary waves when $\mu > \mu_0$, but these solitary waves do not exist when $\mu < \mu_0$. On the other hand, if $\langle u_0, \psi \rangle$ and $\langle G_2, \psi^3 \rangle$ have the opposite sign, b_1^{\pm} are purely imaginary. In this case, the perturbation series (15) give two real-valued solitary waves when $\mu < \mu_0$ but not when $\mu > \mu_0$.

The above perturbation calculations can be continued to higher orders. We can show that the two real solitary-wave solutions (15), which exist on only one side of $\mu = \mu_0$, can be constructed to all orders of $(\mu - \mu_0)^{1/2}$. In addition, these two solitary waves $u^{\pm}(\mathbf{x}; \mu)$ merge with each other when $\mu \rightarrow \mu_0$. We can also show that except these two solitary-wave branches, there are no other solitary-wave solutions near the bifurcation point. Thus a saddle-node bifurcation occurs at $\mu = \mu_0$. This completes the proof of Theorem 1.

Stability properties of solitary waves near saddle-node bifurcations is an important issue. In finite-dimensional dynamical systems, the stability of fixed points always switches at saddle-node bifurcations, and this switching is caused by a linear-stability eigenvalue of the fixed points crossing zero along the real axis [1]. For solitary waves in nonlinear partial differential equations (which can be viewed as fixed points in infinite-dimensional dynamical systems), it is widely believed that their stability also always switches at saddle-node bifurcations. We find that this belief is incorrect. Below, we show that for solitary waves (2) in the generalized NLS equations (1), there are no linear-stability eigenvalues crossing zero at a saddle-node bifurcation point, thus stability switching does not occur.

To study the linear stability of solitary waves (2) in Eq. (1), we perturb them as [12]

$$U(\mathbf{x}, t) = e^{i\mu t} \{u(\mathbf{x}) + [v(\mathbf{x}) + w(\mathbf{x})]e^{\lambda t} + [v^*(\mathbf{x}) - w^*(\mathbf{x})]e^{\lambda^* t}\}, \quad (16)$$

where $v, w \ll 1$ are normal-mode perturbations, and λ is the mode's eigenvalue. Inserting this perturbed solution into Eq. (1) and linearizing, we obtain the following linear-stability eigenvalue problem:

$$\mathcal{L}\Phi = -i\lambda\Phi, \quad (17)$$

where

$$\mathcal{L} = \begin{bmatrix} 0 & L_0 \\ L_1 & 0 \end{bmatrix}, \quad \Phi = \begin{bmatrix} v \\ w \end{bmatrix}, \quad (18)$$

$$L_0 = \nabla^2 - \mu + F(u^2, \mathbf{x}), \quad (19)$$

and L_1 has been given in Eq. (4).

At a saddle-node bifurcation point $\mu = \mu_0$, we denote

$$L_{00} = L_0|_{\mu=\mu_0, u=u_0}, \quad \mathcal{L}_0 = \mathcal{L}|_{\mu=\mu_0, u=u_0}. \quad (20)$$

Then in view of Eq. (3), we have

$$L_{00} u_0 = 0, \quad (21)$$

thus zero is a discrete eigenvalue of L_{00} . From this equation as well as Eq. (6), we have

$$\mathcal{L}_0 \begin{bmatrix} 0 \\ u_0 \end{bmatrix} = \mathcal{L}_0 \begin{bmatrix} \psi \\ 0 \end{bmatrix} = 0, \quad (22)$$

thus zero is also a discrete eigenvalue of \mathcal{L}_0 . On the bifurcation of the zero eigenvalue in \mathcal{L}_0 when μ moves away from μ_0 , we have the following main result.

Theorem 2. Assuming that zero is a simple discrete eigenvalue of L_{00} and L_{10} , then at a saddle-node bifurcation point μ_0 , no eigenvalues of the linear-stability operator \mathcal{L} cross zero, thus no stability switching occurs.

We will give two different proofs for this theorem.

First proof. The idea of this proof is to show that, when μ moves away from μ_0 , the algebraic multiplicity of the zero eigenvalue in \mathcal{L} does not decrease, thus the zero eigenvalue in \mathcal{L} cannot bifurcate out to nonzero.

At the saddle-node bifurcation point $\mu = \mu_0$, $(0, u_0)^T$ and $(\psi, 0)^T$ are two linearly independent eigenfunctions of the zero eigenvalue in \mathcal{L}_0 in view of Eq. (22). Here the superscript “ T ”

represents the transpose of a vector. Under the assumption of Theorem 2, zero is a simple discrete eigenvalue of L_{00} and L_{10} . Thus it is easy to see that \mathcal{L}_0 does not admit any additional eigenfunctions at the zero eigenvalue, which means that the geometric multiplicity of the zero eigenvalue in \mathcal{L}_0 is 2. To determine the algebraic multiplicity of the zero eigenvalue in \mathcal{L}_0 , we need to examine the number of generalized eigenfunctions of this zero eigenvalue. The lowest-order generalized eigenfunction $(f_1, g_1)^T$ to the eigenfunction $(0, u_0)^T$ of this zero eigenvalue satisfies the equation

$$\mathcal{L}_0 \begin{bmatrix} f_1 \\ g_1 \end{bmatrix} = \begin{bmatrix} 0 \\ u_0 \end{bmatrix}, \quad (23)$$

so the equation for f_1 is

$$L_{10} f_1 = u_0. \quad (24)$$

From Eq. (6), we see that this inhomogeneous equation has a homogeneous localized solution ψ . In addition, from conditions of saddle-node bifurcations in Theorem 1, $\langle u_0, \psi \rangle \neq 0$. Furthermore, L_{10} is a self-adjoint operator. Thus, from the Fredholm alternative theorem, the inhomogeneous equation (24) does not admit any localized solution, which means that the eigenfunction $(0, u_0)^T$ of the zero eigenvalue in \mathcal{L}_0 does not have any generalized eigenfunctions. Similarly, we can show that the eigenfunction $(\psi, 0)^T$ of the zero eigenvalue in \mathcal{L}_0 does not have any generalized eigenfunctions either. Hence the algebraic multiplicity of the zero eigenvalue in \mathcal{L}_0 is equal to its geometric multiplicity and is 2.

Away from the bifurcation point (i.e., $\mu \neq \mu_0$), \mathcal{L} always has a zero eigenmode,

$$\mathcal{L} \begin{bmatrix} 0 \\ u \end{bmatrix} = 0, \quad (25)$$

in view of Eq. (3). In addition, by differentiating Eq. (3) with respect to μ , we also get

$$\mathcal{L} \begin{bmatrix} u_\mu \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ u \end{bmatrix}, \quad (26)$$

thus $(u_\mu, 0)^T$ is a generalized eigenfunction of the zero eigenvalue in \mathcal{L} . This means that the algebraic multiplicity of the zero eigenvalue in \mathcal{L} is at least 2 when $\mu \neq \mu_0$.

If nonzero eigenvalues bifurcate out from the zero eigenvalue in \mathcal{L} , the algebraic multiplicity of this zero eigenvalue must decrease. Our results above show that, when μ moves away from μ_0 , the algebraic multiplicity of the zero eigenvalue in \mathcal{L} does not decrease, thus there cannot be nonzero eigenvalues of \mathcal{L} bifurcating out from zero. Consequently, no eigenvalues of \mathcal{L} cross zero at the saddle-node bifurcation point, thus no stability switching occurs. This completes the proof of Theorem 2.

Second proof. In this proof, we directly show nonexistence of nonzero eigenvalues near the bifurcation point. First, by taking the inner products of equations (17) with u and u_μ , we find that for nonzero eigenvalues λ , eigenfunctions (v, w) satisfy the following orthogonality constraints [12]:

$$\langle u, v \rangle = \langle u_\mu, w \rangle = 0. \quad (27)$$

Now suppose at a saddle-node bifurcation point μ_0 , nonzero eigenvalues λ bifurcate out from the origin, i.e., $\lambda \rightarrow 0$ as

$\mu \rightarrow \mu_0$. Then in view of Eq. (17) and our assumption in Theorem 2,

$$(v, w) \rightarrow (c_1 \psi, c_2 u_0), \quad \text{as } \mu \rightarrow \mu_0, \quad (28)$$

where c_1 and c_2 are certain constants which cannot be zero simultaneously. In addition, we see from Eq. (15) that when $\mu \rightarrow \mu_0$, $u_\mu \propto \psi$. Upon substituting these expressions into the orthogonality conditions (27) and taking the limit of $\mu \rightarrow \mu_0$, we find that

$$c_1 \langle u_0, \psi \rangle = c_2 \langle \psi, u_0 \rangle = 0. \quad (29)$$

Since at a saddle-node bifurcation, $\langle u_0, \psi \rangle \neq 0$ (see Theorem 1), the above relations then give $c_1 = c_2 = 0$, which contradicts our earlier requirement of c_1 and c_2 not being zero simultaneously. Thus there cannot be eigenvalues bifurcating out from the origin at a saddle-node bifurcation. This also proves Theorem 2.

Now we discuss the case when Eq. (1) is nonconservative, i.e., the function $F(\dots)$ in Eq. (1) is complex valued. In this case, if F admits parity-time (PT) symmetry $F^*(|U|^2, \mathbf{x}) = F(|U|^2, -\mathbf{x})$, then solitary waves (2) can still exist over a continuous range of real μ values [13], and saddle-node bifurcations can also occur (see later text). By slightly modifying the analysis above, we can show that there is no stability switching at saddle-node bifurcations in these nonconservative systems either. Next we use two examples to confirm the above analytical findings.

Example 1. Consider Eq. (1) with a symmetric double-well potential and cubic-quintic nonlinearity, i.e.,

$$iU_t + U_{xx} - V(x)U + |U|^2U - |U|^4U = 0, \quad (30)$$

where the double-well potential

$$V(x) = -3[\text{sech}^2(x + 1.5) + \text{sech}^2(x - 1.5)] \quad (31)$$

is shown in Fig. 1(a). We have computed solitary waves (2) in this system by the Newton-conjugate-gradient method [12], and their power curve is plotted in Fig. 1(b). Here the soliton power P is defined as $\int_{-\infty}^{\infty} |u|^2 dx$. It is seen that a saddle-node bifurcation occurs at $\mu_0 \approx 2.16$. Two solitary waves on




FIG. 1. (Color online) (a) Potential (31) in Example 1; (b) power curve of solitons; (c), (d) soliton profiles at points marked by the same letters in (b); (e), (f) stability spectra of solitons in (c) and (d).




FIG. 2. (Color online) (a) Power curve of solitons in Example 2; (b) soliton profile at point “b” in (a) (solid and dashed lines for real and imaginary parts); (c),(d) stability spectra of solitons at points “b” and “d” in (a).

the lower and upper branches near this bifurcation point are displayed in Figs. 1(c) and 1(d).

To determine the linear stability of these solitary waves, we have computed their whole linear-stability spectra by the Fourier collocation method [12]. These spectra for the two solitary waves in Figs. 1(c) and 1(d) are shown in Figs. 1(e) and 1(f), respectively. It is seen that none of the spectra contain unstable eigenvalues, indicating that these solitary waves on both lower and upper branches are linearly stable. We have also performed this spectrum computation for other solitary waves on the power curve of Fig. 1(b), and found that they are all linearly stable. Thus there is no stability switching at the saddle-node bifurcation point. Additionally, we have found that the zero eigenvalue for all these solitons has algebraic multiplicity 2, in agreement with our analytical result.

Example 2. We still consider Eq. (30) but now with a complex PT-symmetric potential

$$V(x) = -3[\operatorname{sech}^2(x + 1.5) + \operatorname{sech}^2(x - 1.5)] + 0.25i[\operatorname{sech}^2(x + 1.5) - \operatorname{sech}^2(x - 1.5)]. \quad (32)$$

This nonconservative system still admits solitary waves (2) for continuous real ranges of μ , but $u(x)$ is complex valued now. We have numerically obtained a family of these solitons, whose power curve is plotted in Fig. 2(a). Again a saddle-node bifurcation can be seen at $\mu_0 \approx 2.02$. For solitary waves on the lower and upper branches near this bifurcation point [see Fig. 2(b)], their stability spectra lie entirely on the imaginary axis [see Figs. 2(c) and 2(d)], indicating that they are all linearly stable. Hence no stability switching occurs at saddle-node bifurcations in this nonconservative system either.

In summary, we have shown that for solitary waves in the generalized NLS equations (1) with real or complex potentials, stability does not switch at saddle-node bifurcations. This disproves a widespread belief that such stability switching should always occur in nonlinear partial differential equations. Since the generalized NLS equations (1) arise frequently in nonlinear optics, Bose-Einstein condensates, and other physical disciplines, our finding could have broad impact.

This work is supported in part by the Air Force Office of Scientific Research (USAF 9550-09-1-0228) and the National Science Foundation (DMS-0908167).

-
- [1] J. Guckenheimer and P. Holmes, *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields* (Springer-Verlag, New York, 1990).
- [2] B. Buffoni, A. R. Champneys, and J. F. Toland, *J. Dyn. Differ. Equ.* **8**, 221 (1996).
- [3] T. S. Yang and T. R. Akylas, *J. Fluid Mech.* **30**, 215 (1997).
- [4] M. Chen, *Appl. Anal.* **75**, 213 (2000).
- [5] J. Burke and E. Knobloch, *Chaos* **17**, 037102 (2007).
- [6] G. Herring *et al.*, *Phys. Lett. A* **345**, 144 (2005).
- [7] T. Kapitula, P. Kevrekidis, and Z. Chen, *SIAM J. Appl. Dyn.* **5**, 598 (2006).
- [8] A. Sacchetti, *Phys. Rev. Lett.* **103**, 194101 (2009).
- [9] T. R. Akylas, G. Hwang, and J. Yang, *Proc. R. Soc. London A* **468**, 116 (2012).
- [10] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, *Rev. Mod. Phys.* **71**, 463 (1999).
- [11] Y. S. Kivshar and G. P. Agrawal, *Optical Solitons* (Academic, San Diego, 2003).
- [12] J. Yang, *Nonlinear Waves in Integrable and Nonintegrable Systems* (SIAM, Philadelphia, 2010).
- [13] C. M. Bender and S. Boettcher, *Phys. Rev. Lett.* **80**, 5243 (1998).