

General high-order rogue waves and their dynamics in the nonlinear Schrödinger equation

BY YASUHIRO OHTA^{1,*} AND JIANKE YANG^{2,*}

¹*Department of Mathematics, Kobe University, Rokko, Kobe 657-8501, Japan*

²*Department of Mathematics and Statistics, University of Vermont,
Burlington, VT 05401, USA*

General high-order rogue waves in the nonlinear Schrödinger equation are derived by the bilinear method. These rogue waves are given in terms of determinants whose matrix elements have simple algebraic expressions. It is shown that the general N -th order rogue waves contain $N - 1$ free irreducible complex parameters. In addition, the specific rogue waves obtained by Akhmediev *et al.* (Akhmediev *et al.* 2009 *Phys. Rev. E* **80**, 026601 (doi:10.1103/PhysRevE.80.026601)) correspond to special choices of these free parameters, and they have the highest peak amplitudes among all rogue waves of the same order. If other values of these free parameters are taken, however, these general rogue waves can exhibit other solution dynamics such as arrays of fundamental rogue waves arising at different times and spatial positions and forming interesting patterns.

Keywords: rogue waves; nonlinear Schrödinger equation; bilinear method

1. Introduction

Rogue waves, also known as freak waves, monster waves, killer waves, extreme waves and abnormal waves, is a hot topic in physics these days. This name comes originally from oceanography, and it refers to large and spontaneous ocean surface waves that occur in the sea and are a threat even to large ships and ocean liners. Recently, an optical analogue of rogue waves—optical rogue waves, was observed in optical fibres (Solli *et al.* 2007; Kibler *et al.* 2010). These optical rogue waves are narrow pulses that emerge from initially weakly modulated continuous-wave signals. A growing consensus is that both oceanic and optical rogue waves appear as a result of modulation instability of monochromatic nonlinear waves. Mathematically, the simplest and most universal model for the description of modulation instability and subsequent nonlinear evolution of quasi-monochromatic waves is the focusing nonlinear Schrödinger (NLS) equation (Benney & Newell 1967; Zakharov 1968; Hasegawa & Tappert 1973). This equation is integrable (Zakharov & Shabat 1972), thus its solutions often admit analytical expressions. For rogue waves, the simplest (lowest-order) analytical solution was obtained by Peregrine (1983). This solution approaches a non-zero constant background as time goes to $\pm\infty$, but rises to a peak amplitude of three

*Authors for correspondence (ohta@math.kobe-u.ac.jp; jyang@math.uvm.edu).

times the background in the intermediate time. Special higher order rogue waves were obtained by Akhmediev *et al.* (2009a) using Darboux transformation. These rogue waves could reach higher peak amplitude from a constant background. Recently, more general higher order (multi-Peregrine) rogue waves were obtained by Dubard *et al.* (2010), Dubard & Matveev (2011), Gaillard (2011), Ankiewicz *et al.* (2011) and Kedziora *et al.* (2011). It was shown that these higher order waves could possess multiple intensity peaks at different points of the space–time plane. These exact rogue-wave solutions, which sit on non-zero constant background, are very different from the familiar soliton and multi-soliton solutions which sit on the zero background. They were little known until recently owing to high public interest in theoretical explanations for freak waves observed in the ocean. These rogue waves are intimately related to homoclinic solutions (Akhmediev *et al.* 1985, 1988; Its *et al.* 1988; Ablowitz & Herbst 1990). Indeed, rogue waves can be obtained from homoclinic solutions when the spatial period of homoclinic solutions goes to infinity (Akhmediev *et al.* 1985, 1988, 2009b; Gaillard 2011). These rogue waves are also related to breather solutions which move on a non-zero constant background with profiles changing with time (Akhmediev *et al.* 2009c).

In this article, we derive general high-order rogue waves in the NLS equation and explore their new solution dynamics. Our derivation is based on the bilinear method in the soliton theory (Hirota 2004). Our solution is given in terms of Gram determinants and then further simplified, so that the elements in the determinant matrices have simple algebraic expressions. Compared with the high-order rogue waves presented in Dubard *et al.* (2010) and Gaillard (2011), our solution appears to be more explicit and more easily yielding specific expressions for rogue waves of any given order. We also show that these general rogue waves of N -th order contain $N - 1$ free irreducible complex parameters. In addition, the specific rogue waves obtained in Akhmediev *et al.* (2009a) correspond to special choices of these free parameters, and they have the highest peak amplitudes among all rogue waves of the same order. If other values of these free parameters are taken, however, these general rogue waves can exhibit other solution dynamics such as arrays of fundamental (Peregrine) rogue waves arising at different times and spatial positions. Interesting patterns of these rogue-wave arrays are also illustrated.

2. General rogue-wave solutions

In this paper, we consider general rogue waves in the focusing NLS equation

$$iu_t = u_{xx} + 2|u|^2u. \quad (2.1)$$

Rogue waves are nonlinear waves which approach a constant background at large time and distances. Notice that equation (2.1) is invariant under scalings $x \rightarrow \alpha x$, $t \rightarrow \alpha^2 t$, $u \rightarrow u/\alpha$ for any real constant α . In addition, it is invariant under the Galilean transformation $u(x, t) \rightarrow u(x - vt, t) \exp(-ivx/2 + iv^2t/4)$ for any real velocity v . Thus, we only consider rogue waves which approach unit-amplitude background at large x and t ,

$$u(x, t) \rightarrow e^{-2it} \quad \text{as} \quad x, t \rightarrow \pm\infty.$$

Then under the variable transform $u \rightarrow ue^{-2it}$, the NLS equation (2.1) becomes

$$iu_t = u_{xx} + 2(|u|^2 - 1)u, \quad (2.2)$$

where

$$u(x, t) \rightarrow 1, \quad x, t \rightarrow \pm\infty. \quad (2.3)$$

The rogue waves are described by rational solutions in the NLS equation. In order to present these solutions, let us introduce the so-called elementary Schur polynomials $S_n(\mathbf{x})$ which are defined via the generating function,

$$\sum_{n=0}^{\infty} S_n(\mathbf{x})\lambda^n = \exp\left(\sum_{k=1}^{\infty} x_k\lambda^k\right),$$

where $\mathbf{x} = (x_1, x_2, \dots)$. For example, we have

$$S_0(\mathbf{x}) = 1, \quad S_1(\mathbf{x}) = x_1, \quad S_2(\mathbf{x}) = \frac{1}{2}x_1^2 + x_2, \quad S_3(\mathbf{x}) = \frac{1}{6}x_1^3 + x_1x_2 + x_3, \dots$$

It is known that the general Schur polynomials give the complete set of homogeneous-weight algebraic solutions for the Kadomtsev–Petviashvili (KP) hierarchy (Sato 1981; Jimbo & Miwa 1983).

Theorem 2.1. *The NLS equation (2.2) under the boundary condition (2.3) has non-singular rational solutions*

$$u = \frac{\sigma_1}{\sigma_0}, \quad (2.4)$$

where

$$\sigma_n = \det_{1 \leq i, j \leq N} (m_{2i-1, 2j-1}^{(n)}), \quad (2.5)$$

the matrix elements in σ_n are defined by

$$\left. \begin{aligned} m_{ij}^{(n)} &= \sum_{\nu=0}^{\min(i,j)} \Phi_{i\nu}^{(n)} \Psi_{j\nu}^{(n)}, & \Phi_{i\nu}^{(n)} &= \frac{1}{2^\nu} \sum_{k=0}^{i-\nu} a_k S_{i-\nu-k}(\mathbf{x}^+(n) + \nu \mathbf{s}) \\ \text{and } \Psi_{j\nu}^{(n)} &= \frac{1}{2^\nu} \sum_{l=0}^{j-\nu} \bar{a}_l S_{j-\nu-l}(\mathbf{x}^-(n) + \nu \mathbf{s}), \end{aligned} \right\} \quad (2.6)$$

a_k ($k=0, 1, \dots$) are complex constants, and $\mathbf{x}^\pm(n) = (x_1^\pm(n), x_2^\pm, \dots)$, $\mathbf{s} = (s_1, s_2, \dots)$ are defined by

$$\left. \begin{aligned} x_1^\pm(n) &= x \mp 2it \pm n - \frac{1}{2}, & x_k^\pm &= \frac{x \mp 2^k it}{k!} - r_k, \quad (k \geq 2), \\ \sum_{k=1}^{\infty} r_k \lambda^k &= \ln\left(\cosh \frac{\lambda}{2}\right) & \text{and } \sum_{k=1}^{\infty} s_k \lambda^k &= \ln\left(\frac{2}{\lambda} \tanh \frac{\lambda}{2}\right). \end{aligned} \right\} \quad (2.7)$$

In equation (2.6), \bar{a}_l is the complex conjugate of a_l . The above σ_n can also be expressed as

$$\sigma_n = \sum_{\nu_1=0}^1 \sum_{\nu_2=\nu_1+1}^3 \sum_{\nu_3=\nu_2+1}^5 \cdots \sum_{\nu_N=\nu_{N-1}+1}^{2N-1} \det_{1 \leq i, j \leq N} (\Phi_{2i-1, \nu_j}^{(n)}) \det_{1 \leq i, j \leq N} (\Psi_{2i-1, \nu_j}^{(n)}), \quad (2.8)$$

where we further define

$$\Phi_{iv}^{(n)} = 0, \quad \Psi_{iv}^{(n)} = 0, \quad (i < v). \quad (2.9)$$

Before deriving these rogue wave solutions in this theorem, we give some comments. In the above definitions of r_k and s_k , since the generators are even functions, all odd terms are zero, i.e. $r_1 = r_3 = r_5 = \dots = 0$ and $s_1 = s_3 = s_5 = \dots = 0$. The even-term coefficients are

$$r_2 = \frac{1}{8}, \quad r_4 = -\frac{1}{192}, \quad r_6 = \frac{1}{2880}, \dots, \quad s_2 = -\frac{1}{12}, \quad s_4 = \frac{7}{1440}, \quad s_6 = -\frac{31}{90720}, \dots$$

In the solutions, a_k are complex parameters. We will show in the appendix that without any loss of generality, we can set

$$a_0 = 1, \quad a_2 = a_4 = \dots = a_{\text{even}} = 0.$$

In addition, by a shift of the x and t axes, we can make $a_1 = 0$. Thus, these solutions have $N - 1$ irreducible complex parameters, $a_3, a_5, \dots, a_{2N-1}$.

3. Derivation of general rogue-wave solutions

In this section, we derive the general rogue-wave solutions given in theorem 2.1. This derivation uses the bilinear method in the soliton theory (Hirota 2004). The outline of this derivation is as follows. The NLS equation (2.2) is first transformed into the bilinear form,

$$\left. \begin{aligned} (D_x^2 + 2)f \cdot f &= 2|g|^2 \\ \text{and } (D_x^2 - iD_t)g \cdot f &= 0, \end{aligned} \right\} \quad (3.1)$$

by the variable transformation

$$u = \frac{g}{f}, \quad (3.2)$$

where f is a real variable and g a complex one. Here, D is the Hirota's bilinear differential operator defined by

$$\begin{aligned} &P(D_x, D_y, D_t, \dots)F(x, y, t, \dots) \cdot G(x, y, t, \dots) \\ &= P(\partial_x - \partial_{x'}, \partial_y - \partial_{y'}, \partial_t - \partial_{t'}, \dots)F(x, y, t, \dots)G(x', y', t', \dots)|_{x'=x, y'=y, t'=t, \dots}, \end{aligned}$$

where P is a polynomial of D_x, D_y, D_t, \dots . Then, we consider a 2 + 1-dimensional generalization of the above bilinear equation,

$$\left. \begin{aligned} (D_x D_y + 2)f \cdot f &= 2gh \\ \text{and } (D_x^2 - iD_t)g \cdot f &= 0, \end{aligned} \right\} \quad (3.3)$$

where h is another complex variable. This is in fact the bilinear form of the Davey–Stewartson equation, which is a $2 + 1$ -dimensional generalization of the NLS equation. We first construct a wide class of solutions for equation (3.3) in the form of Gram determinants. If the solutions f , g and h of equation (3.3) further satisfy the conditions,

$$(\partial_x - \partial_y)f = Cf \quad (3.4)$$

and

$$f : \text{real}, \quad h = \bar{g}, \quad (3.5)$$

where C is a constant and the overbar $\bar{}$ represents complex conjugation, then these solutions also satisfy the bilinear NLS equation (3.1). Among the determinant solutions for the $2 + 1$ -dimensional system (3.3), we extract algebraic solutions satisfying the reduction condition (3.4). Then such algebraic solutions satisfy both (3.3) and (3.4), i.e. they are solutions for the $1 + 1$ -dimensional system,

$$\left. \begin{aligned} (D_x^2 + 2)f \cdot f &= 2gh \\ \text{and } (D_x^2 - iD_t)g \cdot f &= 0. \end{aligned} \right\} \quad (3.6)$$

Finally, we impose the real and complex conjugate condition (3.5) on the algebraic solutions. Then the bilinear system (3.6) reduces to the bilinear NLS equation (3.1), hence equation (3.2) gives the general high-order rogue-wave solutions for the NLS equation (2.2).

The execution of the above derivation will involve some novel techniques which are uncommon in the bilinear solution method (Hirota 2004). It is known that the bilinear equations of the NLS hierarchy admit homogeneous-weight polynomial solutions given by the Schur polynomials associated with rectangular Young diagrams (Ikeda & Yamada 2002). However, those solutions do not satisfy the complex conjugation condition $h = \bar{g}$ in general, since the Schur polynomials g and h in Ikeda & Yamada (2002) have different degrees unless the Young diagram associated with f is a square. In the case of a square-shape Young diagram for f , h can be equal to $-\bar{g}$ (but not \bar{g}) and the equation becomes the defocusing NLS equation. To construct rational solutions for the focusing NLS equation (3.1), it is crucial to consider weight-inhomogeneous polynomials. In order to satisfy the reduction condition (3.4) as well as the complex conjugate condition (3.5), we need a specific combination of Schur polynomials as given in theorem 2.1.

Next, we follow the above outline to derive general rogue-wave solutions to the NLS equation (2.2) under the boundary condition (2.3).

(a) Gram determinant solution for the $2 + 1$ -dimensional system

In this subsection, we first derive the Gram determinant solution for the $2 + 1$ -dimensional bilinear equations (3.3).

Lemma 3.1. Let $m_{ij}^{(n)}$, $\varphi_i^{(n)}$ and $\psi_j^{(n)}$ be functions of x_1 , x_2 and x_{-1} satisfying the following differential and difference relations,

$$\left. \begin{aligned} \partial_{x_1} m_{ij}^{(n)} &= \varphi_i^{(n)} \psi_j^{(n)}, \\ \partial_{x_2} m_{ij}^{(n)} &= \varphi_i^{(n+1)} \psi_j^{(n)} + \varphi_i^{(n)} \psi_j^{(n-1)}, \\ \partial_{x_{-1}} m_{ij}^{(n)} &= -\varphi_i^{(n-1)} \psi_j^{(n+1)}, \\ m_{ij}^{(n+1)} &= m_{ij}^{(n)} + \varphi_i^{(n)} \psi_j^{(n+1)} \\ \text{and} \quad \partial_{x_k} \varphi_i^{(n)} &= \varphi_i^{(n+k)}, \quad \partial_{x_k} \psi_j^{(n)} = -\psi_j^{(n-k)}, \quad (k = 1, 2, -1). \end{aligned} \right\} \quad (3.7)$$

Then the determinant,

$$\tau_n = \det_{1 \leq i, j \leq N} (m_{ij}^{(n)}), \quad (3.8)$$

satisfies the bilinear equations,

$$\left. \begin{aligned} (D_{x_1} D_{x_{-1}} - 2)\tau_n \cdot \tau_n &= -2\tau_{n+1}\tau_{n-1} \\ \text{and} \quad (D_{x_1}^2 - D_{x_2})\tau_{n+1} \cdot \tau_n &= 0. \end{aligned} \right\} \quad (3.9)$$

Proof. We have the differential formula of determinant,

$$\partial_x \det_{1 \leq i, j \leq N} (a_{ij}) = \sum_{i, j=1}^N \Delta_{ij} \partial_x a_{ij}, \quad (3.10)$$

and the expansion formula of bordered determinant,

$$\det \begin{pmatrix} a_{ij} & b_i \\ c_j & d \end{pmatrix} = - \sum_{i, j} \Delta_{ij} b_i c_j + d \det(a_{ij}),$$

where Δ_{ij} is the (i, j) -cofactor of the matrix (a_{ij}) . By using these formulae repeatedly, we can verify that the derivatives and shifts of the τ function (3.8) are expressed by the bordered determinants as follows:

$$\begin{aligned} \partial_{x_1} \tau_n &= \begin{vmatrix} m_{ij}^{(n)} & \varphi_i^{(n)} \\ -\psi_j^{(n)} & 0 \end{vmatrix}, \\ \partial_{x_1}^2 \tau_n &= \begin{vmatrix} m_{ij}^{(n)} & \varphi_i^{(n+1)} \\ -\psi_j^{(n)} & 0 \end{vmatrix} + \begin{vmatrix} m_{ij}^{(n)} & \varphi_i^{(n)} \\ \psi_j^{(n-1)} & 0 \end{vmatrix}, \\ \partial_{x_2} \tau_n &= \begin{vmatrix} m_{ij}^{(n)} & \varphi_i^{(n+1)} \\ -\psi_j^{(n)} & 0 \end{vmatrix} - \begin{vmatrix} m_{ij}^{(n)} & \varphi_i^{(n)} \\ \psi_j^{(n-1)} & 0 \end{vmatrix}, \end{aligned}$$

$$\begin{aligned} \partial_{x_{-1}} \tau_n &= \begin{vmatrix} m_{ij}^{(n)} & \varphi_i^{(n-1)} \\ \psi_j^{(n+1)} & 0 \end{vmatrix}, \\ (\partial_{x_1} \partial_{x_{-1}} - 1) \tau_n &= \begin{vmatrix} m_{ij}^{(n)} & \varphi_i^{(n-1)} & \varphi_i^{(n)} \\ \psi_j^{(n+1)} & 0 & -1 \\ -\psi_j^{(n)} & -1 & 0 \end{vmatrix}, \\ \tau_{n+1} &= \begin{vmatrix} m_{ij}^{(n)} & \varphi_i^{(n)} \\ -\psi_j^{(n+1)} & 1 \end{vmatrix}, \\ \tau_{n-1} &= \begin{vmatrix} m_{ij}^{(n)} & \varphi_i^{(n-1)} \\ \psi_j^{(n)} & 1 \end{vmatrix}, \\ \partial_{x_1} \tau_{n+1} &= \begin{vmatrix} m_{ij}^{(n)} & \varphi_i^{(n+1)} \\ -\psi_j^{(n+1)} & 0 \end{vmatrix}, \\ \partial_{x_1}^2 \tau_{n+1} &= \begin{vmatrix} m_{ij}^{(n)} & \varphi_i^{(n+2)} \\ -\psi_j^{(n+1)} & 0 \end{vmatrix} + \begin{vmatrix} m_{ij}^{(n)} & \varphi_i^{(n)} & \varphi_i^{(n+1)} \\ -\psi_j^{(n)} & 0 & 0 \\ -\psi_j^{(n+1)} & 1 & 0 \end{vmatrix}, \\ \text{and } \partial_{x_2} \tau_{n+1} &= \begin{vmatrix} m_{ij}^{(n)} & \varphi_i^{(n+2)} \\ -\psi_j^{(n+1)} & 0 \end{vmatrix} - \begin{vmatrix} m_{ij}^{(n)} & \varphi_i^{(n)} & \varphi_i^{(n+1)} \\ -\psi_j^{(n)} & 0 & 0 \\ -\psi_j^{(n+1)} & 1 & 0 \end{vmatrix}. \end{aligned}$$

From the Jacobi formula of determinants,

$$\begin{vmatrix} a_{ij} & b_i & c_i \\ d_j & e & f \\ g_j & h & k \end{vmatrix} \times |a_{ij}| = \begin{vmatrix} a_{ij} & c_i \\ g_j & k \end{vmatrix} \times \begin{vmatrix} a_{ij} & b_i \\ d_j & e \end{vmatrix} - \begin{vmatrix} a_{ij} & b_i \\ g_j & h \end{vmatrix} \times \begin{vmatrix} a_{ij} & c_i \\ d_j & f \end{vmatrix},$$

we immediately obtain the identities,

$$\begin{aligned} (\partial_{x_1} \partial_{x_{-1}} - 1) \tau_n \times \tau_n &= \partial_{x_1} \tau_n \times \partial_{x_{-1}} \tau_n - (-\tau_{n-1})(-\tau_{n+1}), \\ \frac{1}{2}(\partial_{x_1}^2 - \partial_{x_2}) \tau_{n+1} \times \tau_n &= \partial_{x_1} \tau_{n+1} \times \partial_{x_1} \tau_n - \tau_{n+1} \frac{1}{2}(\partial_{x_1}^2 + \partial_{x_2}) \tau_n, \end{aligned}$$

which are the bilinear equations (3.9). This completes the proof. ■

Since the matrix element $m_{ij}^{(n)}$ is written as

$$m_{ij}^{(n)} = \int^{x_1} \varphi_i^{(n)} \psi_j^{(n)} dx_1,$$

the determinant (3.8) is often called the Gram determinant solution. Let us define

$$f = \tau_0, \quad g = \tau_1, \quad h = \tau_{-1},$$

then these are the Gram determinant solution for the 2 + 1-dimensional system,

$$(D_{x_1} D_{x_{-1}} - 2)f \cdot f = -2gh$$

and

$$(D_{x_1}^2 - D_{x_2})g \cdot f = 0,$$

which is nothing but the bilinear equations (3.3) by writing $x_1 = x$, $x_2 = -it$ and $x_{-1} = -y$.

(b) *Algebraic solutions for the 1 + 1-dimensional system*

Next, we derive algebraic solutions satisfying both the bilinear equations (3.3) and the reduction condition (3.4), hence satisfying the 1 + 1-dimensional system (3.6). These solutions are obtained by choosing the matrix elements appropriately in the Gram determinant solution in lemma 3.1.

Lemma 3.2. *We define matrix elements $m_{ij}^{(n)}$ by*

$$m_{ij}^{(n)} = A_i B_j m^{(n)}|_{p=1, q=1} \quad (3.11)$$

and

$$m^{(n)} = \frac{1}{p+q} \left(-\frac{p}{q}\right)^n e^{\xi+\eta}, \quad \xi = px_1 + p^2x_2, \quad \eta = qx_1 - q^2x_2, \quad (3.12)$$

where A_i and B_j are differential operators with respect to p and q , respectively, defined as

$$\begin{aligned} A_0 &= a_0, \\ A_1 &= a_0 p \partial_p + a_1, \\ A_2 &= \frac{a_0}{2} (p \partial_p)^2 + a_1 p \partial_p + a_2, \\ &\vdots \\ A_i &= \sum_{k=0}^i \frac{a_k}{(i-k)!} (p \partial_p)^{i-k}, \end{aligned}$$

and

$$\begin{aligned} B_0 &= b_0, \\ B_1 &= b_0 q \partial_q + b_1, \\ B_2 &= \frac{b_0}{2} (q \partial_q)^2 + b_1 q \partial_q + b_2, \\ &\vdots \\ B_j &= \sum_{l=0}^j \frac{b_l}{(j-l)!} (q \partial_q)^{j-l}, \end{aligned}$$

and a_k and b_l are constants. Then the determinant

$$\tau_n = \det_{1 \leq i, j \leq N} (m_{2i-1, 2j-1}^{(n)}) = \begin{vmatrix} m_{11}^{(n)} & m_{13}^{(n)} & \cdots & m_{1, 2N-1}^{(n)} \\ m_{31}^{(n)} & m_{33}^{(n)} & \cdots & m_{3, 2N-1}^{(n)} \\ \vdots & \vdots & & \vdots \\ m_{2N-1, 1}^{(n)} & m_{2N-1, 3}^{(n)} & \cdots & m_{2N-1, 2N-1}^{(n)} \end{vmatrix} \quad (3.13)$$

satisfies the bilinear equations

$$\left. \begin{aligned} (D_{x_1}^2 + 2)\tau_n \cdot \tau_n &= 2\tau_{n+1}\tau_{n-1} \\ \text{and } (D_{x_1}^2 - D_{x_2})\tau_{n+1} \cdot \tau_n &= 0. \end{aligned} \right\} \quad (3.14)$$

Proof. First let us introduce $\tilde{m}^{(n)}$, $\tilde{\varphi}^{(n)}$ and $\tilde{\psi}^{(n)}$ by

$$\tilde{m}^{(n)} = \frac{1}{p+q} \left(-\frac{p}{q} \right)^n e^{\tilde{\xi} + \tilde{\eta}}, \quad \tilde{\varphi}^{(n)} = p^n e^{\tilde{\xi}}, \quad \tilde{\psi}^{(n)} = (-q)^{-n} e^{\tilde{\eta}},$$

where

$$\tilde{\xi} = \frac{1}{p}x_{-1} + px_1 + p^2x_2 \quad \text{and} \quad \tilde{\eta} = \frac{1}{q}x_{-1} + qx_1 - q^2x_2.$$

Obviously, these functions satisfy the differential and difference relations

$$\begin{aligned} \partial_{x_1} \tilde{m}^{(n)} &= \tilde{\varphi}^{(n)} \tilde{\psi}^{(n)}, \\ \partial_{x_2} \tilde{m}^{(n)} &= \tilde{\varphi}^{(n+1)} \tilde{\psi}^{(n)} + \tilde{\varphi}^{(n)} \tilde{\psi}^{(n-1)}, \\ \partial_{x_{-1}} \tilde{m}^{(n)} &= -\tilde{\varphi}^{(n-1)} \tilde{\psi}^{(n+1)}, \\ \tilde{m}^{(n+1)} &= \tilde{m}^{(n)} + \tilde{\varphi}^{(n)} \tilde{\psi}^{(n+1)} \end{aligned}$$

and
$$\partial_{x_k} \tilde{\varphi}^{(n)} = \tilde{\varphi}^{(n+k)}, \quad \partial_{x_k} \tilde{\psi}^{(n)} = -\tilde{\psi}^{(n-k)}, \quad (k = 1, 2, -1).$$

Therefore, by defining

$$\tilde{m}_{ij}^{(n)} = A_i B_j \tilde{m}^{(n)}, \quad \tilde{\varphi}_i^{(n)} = A_i \tilde{\varphi}^{(n)}, \quad \tilde{\psi}_j^{(n)} = B_j \tilde{\psi}^{(n)},$$

we see that these $\tilde{m}_{ij}^{(n)}$, $\tilde{\varphi}_i^{(n)}$ and $\tilde{\psi}_j^{(n)}$ obey the differential and difference relations (3.7) since the operators A_i and B_j commute with differentials ∂_{x_k} . Lemma 3.1 then tells us that for an arbitrary sequence of indices $(i_1, i_2, \dots, i_N; j_1, j_2, \dots, j_N)$, the determinant

$$\tilde{\tau}_n = \det_{1 \leq \nu, \mu \leq N} (\tilde{m}_{i_\nu, j_\mu}^{(n)})$$

satisfies the bilinear equations (3.9). For example,

$$\tilde{\tau}_n = \det_{1 \leq i, j \leq N} (\tilde{m}_{2i-1, 2j-1}^{(n)}),$$

is a solution to equation (3.9), where p and q are arbitrary parameters.

Next, we consider the reduction condition. From the Leibniz rule, we have the operator relation,

$$(p\partial_p)^m \left(p + \frac{1}{p} \right) = \sum_{l=0}^m \binom{m}{l} \left(p + (-1)^l \frac{1}{p} \right) (p\partial_p)^{m-l},$$

thus we get

$$\begin{aligned} A_i \left(p + \frac{1}{p} \right) &= \sum_{k=0}^i \frac{a_k}{(i-k)!} \sum_{l=0}^{i-k} \binom{i-k}{l} \left(p + (-1)^l \frac{1}{p} \right) (p\partial_p)^{i-k-l} \\ &= \sum_{l=0}^i \sum_{k=0}^{i-l} \frac{a_k}{l!(i-k-l)!} \left(p + (-1)^l \frac{1}{p} \right) (p\partial_p)^{i-k-l} \\ &= \sum_{l=0}^i \frac{1}{l!} \left(p + (-1)^l \frac{1}{p} \right) A_{i-l}, \end{aligned}$$

and similarly

$$B_j \left(q + \frac{1}{q} \right) = \sum_{l=0}^j \frac{1}{l!} \left(q + (-1)^l \frac{1}{q} \right) B_{j-l}.$$

By using these relations, we find that $\tilde{m}_{ij}^{(n)}$ satisfies

$$\begin{aligned} (\partial_{x_1} + \partial_{x_{-1}}) \tilde{m}_{ij}^{(n)} &= A_i B_j (\partial_{x_1} + \partial_{x_{-1}}) \tilde{m}^{(n)} = A_i B_j \left(p + q + \frac{1}{p} + \frac{1}{q} \right) \tilde{m}^{(n)} \\ &= \sum_{k=0}^i \frac{1}{k!} \left(p + (-1)^k \frac{1}{p} \right) A_{i-k} B_j \tilde{m}^{(n)} \\ &\quad + \sum_{l=0}^j \frac{1}{l!} \left(q + (-1)^l \frac{1}{q} \right) A_i B_{j-l} \tilde{m}^{(n)} \\ &= \sum_{k=0}^i \frac{1}{k!} \left(p + (-1)^k \frac{1}{p} \right) \tilde{m}_{i-k,j}^{(n)} + \sum_{l=0}^j \frac{1}{l!} \left(q + (-1)^l \frac{1}{q} \right) \tilde{m}_{i,j-l}^{(n)}. \end{aligned}$$

Now let us take $p = 1$ and $q = 1$. Then $\tilde{m}_{ij}^{(n)}|_{p=1,q=1}$ satisfies the contiguity relation,

$$(\partial_{x_1} + \partial_{x_{-1}}) (\tilde{m}_{ij}^{(n)}|_{p=1,q=1}) = 2 \sum_{\substack{k=0 \\ k:\text{even}}}^i \frac{1}{k!} \tilde{m}_{i-k,j}^{(n)} \Big|_{p=1,q=1} + 2 \sum_{\substack{l=0 \\ l:\text{even}}}^j \frac{1}{l!} \tilde{m}_{i,j-l}^{(n)} \Big|_{p=1,q=1}. \quad (3.15)$$

By using the formula (3.10) and the above relation, the differential of the determinant,

$$\tilde{\tau}_n = \det_{1 \leq i,j \leq N} (\tilde{m}_{2i-1,2j-1}^{(n)}|_{p=1,q=1})$$

is calculated as

$$\begin{aligned}
& (\partial_{x_1} + \partial_{x_{-1}}) \tilde{\tau}_n \\
&= \sum_{i=1}^N \sum_{j=1}^N \Delta_{ij} (\partial_{x_1} + \partial_{x_{-1}}) (\tilde{m}_{2i-1, 2j-1}^{(n)} |_{p=1, q=1}) \\
&= \sum_{i=1}^N \sum_{j=1}^N \Delta_{ij} \left(2 \sum_{\substack{k=0 \\ k:\text{even}}}^{2i-1} \frac{1}{k!} \tilde{m}_{2i-1-k, 2j-1}^{(n)} \Big|_{p=1, q=1} + 2 \sum_{\substack{l=0 \\ l:\text{even}}}^{2j-1} \frac{1}{l!} \tilde{m}_{2i-1, 2j-1-l}^{(n)} \Big|_{p=1, q=1} \right) \\
&= 2 \sum_{i=1}^N \sum_{\substack{k=0 \\ k:\text{even}}}^{2i-1} \frac{1}{k!} \sum_{j=1}^N \Delta_{ij} \tilde{m}_{2i-1-k, 2j-1}^{(n)} \Big|_{p=1, q=1} \\
&\quad + 2 \sum_{j=1}^N \sum_{\substack{l=0 \\ l:\text{even}}}^{2j-1} \frac{1}{l!} \sum_{i=1}^N \Delta_{ij} \tilde{m}_{2i-1, 2j-1-l}^{(n)} \Big|_{p=1, q=1},
\end{aligned}$$

where Δ_{ij} is the (i, j) -cofactor of $\text{mat}_{1 \leq i, j \leq N} (\tilde{m}_{2i-1, 2j-1}^{(n)} |_{p=1, q=1})$. In the first term of the right-hand side, only the term with $k=0$ survives and the other terms vanish, since for $k=2, 4, \dots$, the summation with respect to j is a determinant with two identical rows. Similarly in the second term, only the term with $l=0$ remains. Thus, the right side of the above equation becomes

$$2 \sum_{i=1}^N \sum_{j=1}^N \Delta_{ij} \tilde{m}_{2i-1, 2j-1}^{(n)} \Big|_{p=1, q=1} + 2 \sum_{j=1}^N \sum_{i=1}^N \Delta_{ij} \tilde{m}_{2i-1, 2j-1}^{(n)} \Big|_{p=1, q=1} = 4N \tilde{\tau}_n.$$

Therefore, $\tilde{\tau}_n$ satisfies the reduction condition

$$(\partial_{x_1} + \partial_{x_{-1}}) \tilde{\tau}_n = 4N \tilde{\tau}_n. \tag{3.16}$$

Since $\tilde{\tau}_n$ is a special case of $\tilde{\tau}_n$, it also satisfies the bilinear equations (3.9) with τ_n replaced by $\tilde{\tau}_n$. From (3.9) and (3.16), we see that $\tilde{\tau}_n$ satisfies the 1 + 1-dimensional bilinear equations

$$(D_{x_1}^2 + 2) \tilde{\tau}_n \cdot \tilde{\tau}_n = 2 \tilde{\tau}_{n+1} \tilde{\tau}_{n-1}$$

and

$$(D_{x_1}^2 - D_{x_2}) \tilde{\tau}_{n+1} \cdot \tilde{\tau}_n = 0,$$

which are the same as equation (3.14). Now we can take $x_{-1} = 0$, then $\tilde{m}_{ij}^{(n)} |_{p=1, q=1}$ and $\tilde{\tau}_n$ reduce to $m_{ij}^{(n)}$ and τ_n in lemma 3.2, and this τ_n satisfies the bilinear equations (3.14). This completes the proof. \blacksquare

The above proof uses the technique of reduction. The reduction is a procedure to derive solutions of a lower dimensional system from those of a higher dimensional one. By using the reduction condition (3.16), the derivative with respect to a variable x_{-1} is replaced by the derivative with respect to another variable x_1 . Then in the solution, x_{-1} is just a parameter to which we can substitute any value (such as zero as we did above). This reduction technique has recently been used to derive general dark–dark solitons in the coupled NLS equations (Ohta *et al.* 2011).

It is remarkable that the determinant expression of the solution (3.13) has a quite unique structure: the indices of matrix elements, which label the degree of polynomial, have the step of 2. This comes from the requirement of the reduction condition, i.e. since the contiguity relation (3.15) relates matrix elements with indices shifted by even numbers, we want such a determinant to satisfy the reduction condition. This type of Gram determinant solutions has not been reported in the literature to the best of the authors' knowledge.

From lemma 3.2, by writing $x_1 = x$ and $x_2 = -it$, we find that $f = \tau_0$, $g = \tau_1$ and $h = \tau_{-1}$ satisfy the 1 + 1-dimensional system (3.6).

(c) *Complex conjugacy and regularity*

Now we consider the complex conjugate condition (3.5) and the regularity (non-singularity) of solutions. This complex conjugate condition now is

$$\tau_0 : \text{real}, \quad \tau_{-1} = \bar{\tau}_1.$$

Since $x_1 = x$ is real and $x_2 = -it$ is pure imaginary in lemma 3.2, the above condition is easily satisfied by taking the parameters a_k and b_k to be complex conjugate to each other,

$$b_k = \bar{a}_k. \quad (3.17)$$

In fact, under the condition (3.17) we have

$$\overline{m_{ij}^{(n)}} = m_{ij}^{(n)} \Big|_{a_k \leftrightarrow b_k, x_2 \leftrightarrow -x_2} = m_{ji}^{(-n)},$$

and therefore

$$\bar{\tau}_n = \tau_{-n}.$$

Under condition (3.17), we can further show that the rational solution $u = g/f = \tau_1/\tau_0$ is non-singular, i.e. τ_0 is non-zero for all (x, t) . To prove it, we notice that $f = \tau_0$ is the determinant of a Hermitian matrix $M = \text{mat}_{1 \leq i, j \leq N} (m_{2i-1, 2j-1}^{(0)})$. For any non-zero column vector $\mathbf{v} = (v_1, v_2, \dots, v_N)^T$ and $\bar{\mathbf{v}}$ being its complex transpose, we have

$$\begin{aligned} \bar{\mathbf{v}} M \mathbf{v} &= \sum_{i,j=1}^N \bar{v}_i m_{2i-1, 2j-1}^{(0)} v_j = \sum_{i,j=1}^N \bar{v}_i v_j A_{2i-1} B_{2j-1} \frac{1}{p+q} e^{\xi+\eta} \Big|_{p=1, q=1} \\ &= \sum_{i,j=1}^N \bar{v}_i v_j A_{2i-1} B_{2j-1} \int_{-\infty}^x e^{\xi+\eta} dx \Big|_{p=1, q=1} \end{aligned}$$

$$\begin{aligned}
&= \int_{-\infty}^x \sum_{i,j=1}^N \bar{v}_i v_j A_{2i-1} B_{2j-1} e^{\xi+\eta} \Big|_{p=1,q=1} dx \\
&= \int_{-\infty}^x \left| \sum_{i=1}^N \bar{v}_i A_{2i-1} e^{\xi} \Big|_{p=1} \right|^2 dx > 0,
\end{aligned}$$

which proves that the Hermitian matrix M is positive definite. Therefore, the denominator $f = \det M > 0$, so the solution u is non-singular.

It is noted that the above proofs of complex conjugate condition and regularity condition are quite easy. This is an advantage of the Gram determinant expression of solutions (when compared with the Wronskian expression).

Summarizing the above results, we obtain the following intermediate theorem on rogue-wave solutions in the NLS equation.

Theorem 3.3. *The NLS equation (2.2) has the non-singular rational solutions,*

$$u = \frac{\tau_1}{\tau_0}, \quad (3.18)$$

where

$$\tau_n = \det_{1 \leq i,j \leq N} (m_{2i-1,2j-1}^{(n)}), \quad (3.19)$$

where the matrix elements are defined by

$$\begin{aligned}
m_{ij}^{(n)} &= \sum_{k=0}^i \sum_{l=0}^j \frac{a_k}{(i-k)!} \frac{\bar{a}_l}{(j-l)!} (p\partial_p)^{i-k} (q\partial_q)^{j-l} \frac{1}{p+q} \\
&\quad \times \left(-\frac{p}{q} \right)^n e^{(p+q)x - (p^2 - q^2)\sqrt{-1}t} \Big|_{p=1,q=1}, \quad (3.20)
\end{aligned}$$

and a_k are complex constants.

(d) Simplification of rogue-wave solutions

Finally, we simplify the rogue-wave solutions in theorem 3.3 and derive the solution formulae given in theorem 2.1. The generator \mathcal{G} of the differential operators $(p\partial_p)^k (q\partial_q)^l$ is given as

$$\mathcal{G} = \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \frac{\kappa^k}{k!} \frac{\lambda^l}{l!} (p\partial_p)^k (q\partial_q)^l = \exp(\kappa p\partial_p + \lambda q\partial_q) = \exp(\kappa \partial_{\ln p} + \lambda \partial_{\ln q}),$$

thus for any function $F(p, q)$, we have

$$\mathcal{G}F(p, q) = F(e^\kappa p, e^\lambda q). \quad (3.21)$$

This relation can also be seen by expanding its right-hand side into Taylor series of (κ, λ) around the point $(0, 0)$. By applying this relation to

$$m^{(n)} = \frac{1}{p+q} \left(-\frac{p}{q} \right)^n \exp((p+q)x - (p^2 - q^2)it),$$

we get

$$\mathcal{G}m^{(n)} = \frac{1}{e^\kappa p + e^\lambda q} \left(-\frac{e^\kappa p}{e^\lambda q} \right)^n \exp((e^\kappa p + e^\lambda q)x - (e^{2\kappa} p^2 - e^{2\lambda} q^2)it),$$

thus

$$\begin{aligned} & \frac{1}{m^{(n)}} \mathcal{G}m^{(n)} \Big|_{p=1, q=1} \\ &= \frac{2}{e^\kappa + e^\lambda} e^{n(\kappa-\lambda)} \exp((e^\kappa + e^\lambda - 2)x - (e^{2\kappa} - e^{2\lambda})it) \\ &= \frac{1}{1 - (e^\kappa - 1)(e^\lambda - 1)/(e^\kappa + 1)(e^\lambda + 1)} \\ & \quad \times \exp\left(n(\kappa - \lambda) + (e^\kappa + e^\lambda - 2)x - (e^{2\kappa} - e^{2\lambda})it - \ln \frac{(e^\kappa + 1)(e^\lambda + 1)}{4} \right). \end{aligned}$$

In the most right-hand side, the exponent is rewritten as

$$\begin{aligned} & n(\kappa - \lambda) + \sum_{k=1}^{\infty} \frac{\kappa^k}{k!} (x - 2^k it) + \sum_{l=1}^{\infty} \frac{\lambda^l}{l!} (x + 2^l it) - \frac{\kappa}{2} - \frac{\lambda}{2} - \ln \left(\cosh \frac{\kappa}{2} \cosh \frac{\lambda}{2} \right) \\ &= \sum_{k=1}^{\infty} x_k^+ \kappa^k + \sum_{l=1}^{\infty} x_l^- \lambda^l, \end{aligned}$$

where x_k^+ and x_l^- are defined in (2.7), and the prefactor is rewritten as

$$\begin{aligned} \sum_{\nu=0}^{\infty} \left(\frac{(e^\kappa - 1)(e^\lambda - 1)}{(e^\kappa + 1)(e^\lambda + 1)} \right)^\nu &= \sum_{\nu=0}^{\infty} \left(\frac{\kappa\lambda}{4} \right)^\nu \exp \left(\nu \ln \left(\frac{4}{\kappa\lambda} \tanh \frac{\kappa}{2} \tanh \frac{\lambda}{2} \right) \right) \\ &= \sum_{\nu=0}^{\infty} \left(\frac{\kappa\lambda}{4} \right)^\nu \exp \left(\nu \sum_{k=1}^{\infty} s_k (\kappa^k + \lambda^k) \right), \end{aligned}$$

where s_k is defined in (2.7). Therefore, we obtain

$$\frac{1}{m^{(n)}} \mathcal{G}m^{(n)} \Big|_{p=1, q=1} = \sum_{\nu=0}^{\infty} \left(\frac{\kappa\lambda}{4} \right)^\nu \exp \left(\sum_{k=1}^{\infty} (x_k^+ + \nu s_k) \kappa^k + \sum_{l=1}^{\infty} (x_l^- + \nu s_l) \lambda^l \right),$$

where O and I are the $N \times N$ zero matrix and $2N \times 2N$ unit matrix, respectively. Applying the Laplace expansion to the above determinant, we get

$$\sigma_n = \sum_{0 \leq \nu_1 < \nu_2 < \dots < \nu_N \leq 2N-1} \begin{vmatrix} \Phi_{1\nu_1}^{(n)} & \Phi_{1\nu_2}^{(n)} & \dots & \Phi_{1\nu_N}^{(n)} \\ \Phi_{3\nu_1}^{(n)} & \Phi_{3\nu_2}^{(n)} & \dots & \Phi_{3\nu_N}^{(n)} \\ \vdots & \vdots & & \vdots \\ \Phi_{2N-1,\nu_1}^{(n)} & \Phi_{2N-1,\nu_2}^{(n)} & \dots & \Phi_{2N-1,\nu_N}^{(n)} \end{vmatrix} \\ \times \begin{vmatrix} \Psi_{1\nu_1}^{(n)} & \Psi_{3\nu_1}^{(n)} & \dots & \Psi_{2N-1,\nu_1}^{(n)} \\ \Psi_{1\nu_2}^{(n)} & \Psi_{3\nu_2}^{(n)} & \dots & \Psi_{2N-1,\nu_2}^{(n)} \\ \vdots & \vdots & & \vdots \\ \Psi_{1\nu_N}^{(n)} & \Psi_{3\nu_N}^{(n)} & \dots & \Psi_{2N-1,\nu_N}^{(n)} \end{vmatrix},$$

and noticing (2.9), the expanded expression (2.8) is obtained. Theorem 2.1 is then proved.

(e) Boundary conditions

In order to show the boundary asymptotics (2.3), let us estimate the degree of polynomials of the denominator and numerator in (2.4). The elementary Schur polynomial $S_k(\mathbf{x})$ has the form $S_k(\mathbf{x}) = (x_1)^k/k! +$ (lower degree terms), where $\mathbf{x} = (x_1, x_2, \dots)$. Thus, the degree of the polynomial $S_k(\mathbf{x}^\pm + \nu \mathbf{s})$ in (x, t) is k and its leading term appears in the monomial $(x_1^\pm)^k/k!$, i.e. the leading term is given by $(x \mp 2it)^k/k!$. Therefore, the degrees of $\Phi_{j\nu}^{(n)}$ and $\Psi_{j\nu}^{(n)}$ are both $j - \nu$, and their leading terms are $a_0(x - 2it)^{j-\nu}/(j - \nu)!2^\nu$ and $\bar{a}_0(x + 2it)^{j-\nu}/(j - \nu)!2^\nu$, respectively. Therefore, both of the degrees of determinants $\det_{1 \leq i, j \leq N}(\Phi_{2i-1,\nu_j}^{(n)})$ and $\det_{1 \leq i, j \leq N}(\Psi_{2i-1,\nu_j}^{(n)})$ are given by $1 + 3 + \dots + (2N - 1) - \nu_1 - \nu_2 - \dots - \nu_N$, and in the expression (2.8), the highest degree term comes from the term of $\nu_1 = 0, \nu_2 = 1, \dots, \nu_N = N - 1$ in the summation. For $\nu_j = j - 1$, we have

$$\det_{1 \leq i, j \leq N}(\Phi_{2i-1, j-1}^{(n)}) \\ = \begin{vmatrix} a_0 x_1^+ & \frac{a_0}{2} & 0 & 0 & 0 & \dots \\ \frac{a_0 (x_1^+)^3}{3!} & \frac{a_0 (x_1^+)^2}{2!2} & \frac{a_0 x_1^+}{2^2} & \frac{a_0}{2^3} & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & & \\ \frac{a_0 (x_1^+)^{2N-1}}{(2N-1)!} & \frac{a_0 (x_1^+)^{2N-2}}{(2N-2)!2} & \frac{a_0 (x_1^+)^{2N-3}}{(2N-3)!2^2} & \frac{a_0 (x_1^+)^{2N-4}}{(2N-4)!2^3} & \dots & \frac{a_0 (x_1^+)^N}{N!2^{N-1}} \end{vmatrix} \\ + \text{(lower degree terms)}$$

$$= \frac{a_0^N (x_1^+)^{N(N+1)/2}}{1!3! \cdots (2N-1)! 2^{N(N-1)/2}}$$

$$\times \begin{vmatrix} 1 & 1 & 0 & 0 & 0 & \cdots \\ 1 & 3 & 3 \cdot 2 & 3 \cdot 2 \cdot 1 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 2N-1 & (2N-1)(2N-2) & (2N-1)(2N-2)(2N-3) & \cdots & (2N-1)(2N-2) \cdots (N+1) \end{vmatrix}$$

+ (lower degree terms).

The above determinant is equal to

$$\det_{1 \leq i, j \leq N} \left(\prod_{\nu=1}^{j-1} (2i - \nu) \right) = \det_{1 \leq i, j \leq N} ((2i-1)^{j-1}),$$

which is the Vandermonde determinant. Thus, we obtain

$$\det_{1 \leq i, j \leq N} (\Phi_{2i-1, j-1}^{(n)}) = \frac{0!1! \cdots (N-1)!}{1!3! \cdots (2N-1)!} a_0^N (x_1^+)^{N(N+1)/2} + (\text{lower degree terms}),$$

and similarly

$$\det_{1 \leq i, j \leq N} (\Psi_{2i-1, j-1}^{(n)}) = \frac{0!1! \cdots (N-1)!}{1!3! \cdots (2N-1)!} \bar{a}_0^N (x_1^-)^{N(N+1)/2} + (\text{lower degree terms}).$$

Consequently, the leading term of σ_n is given by

$$\left(\frac{0!1! \cdots (N-1)!}{1!3! \cdots (2N-1)!} \right)^2 |a_0|^{2N} (x^2 + 4t^2)^{N(N+1)/2},$$

which is independent of n . Hence $u = \sigma_1/\sigma_0$ satisfies the boundary condition (2.3).

4. Solution dynamics

In this section, we discuss the dynamics of these general rogue-wave solutions.

To obtain the first-order rogue wave, we set $N = 1$ in theorem 2.1. In this case,

$$m_{11}^{(0)} = \left(x - 2it - \frac{1}{2} + a_1\right) \left(x + 2it - \frac{1}{2} + \bar{a}_1\right) + \frac{1}{4}$$

and

$$m_{11}^{(1)} = \left(x - 2it + \frac{1}{2} + a_1\right) \left(x + 2it - \frac{3}{2} + \bar{a}_1\right) + \frac{1}{4},$$

hence the first-order rogue wave is

$$u(x, t) = \frac{m_{11}^{(1)}}{m_{11}^{(0)}} = \frac{\left(x - 2it + \frac{1}{2} + a_1\right) \left(x + 2it - \frac{3}{2} + \bar{a}_1\right) + \frac{1}{4}}{\left(x - 2it - \frac{1}{2} + a_1\right) \left(x + 2it - \frac{1}{2} + \bar{a}_1\right) + \frac{1}{4}}. \quad (4.1)$$

Clearly, the complex parameter a_1 in this solution can be normalized to zero by a shift of x and t , as we have mentioned before. After setting $a_1 = 0$, this first-order

rogue wave can be rewritten as

$$u(x, t) = 1 - \frac{4(1 - 4it)}{1 + 4\hat{x}^2 + 16t^2}, \quad (4.2)$$

where $\hat{x} = x - 1/2$. This rogue wave was first obtained by Peregrine (1983), see also Akhmediev *et al.* (2009a). Its maximum peak amplitude is equal to 3, i.e. three times the background amplitude.

To obtain the second-order rogue waves, we take $N = 2$. In this case,

$$u = \frac{\begin{vmatrix} m_{11}^{(1)} & m_{13}^{(1)} \\ m_{31}^{(1)} & m_{33}^{(1)} \end{vmatrix}}{\begin{vmatrix} m_{11}^{(0)} & m_{13}^{(0)} \\ m_{31}^{(0)} & m_{33}^{(0)} \end{vmatrix}}. \quad (4.3)$$

From the previous discussions, we will set $a_1 = a_2 = 0$. Then the general second-order rogue wave can be obtained from (4.3) as

$$u = 1 + \frac{\phi}{\psi}, \quad (4.4)$$

where

$$\begin{aligned} \phi = & 24\{(3x - 6x^2 + 4x^3 - 2x^4 - 48t^2 + 48xt^2 - 48x^2t^2 - 160t^4) \\ & + it(-12 + 12x - 16x^3 + 8x^4 + 32t^2 - 64xt^2 + 64x^2t^2 + 128t^4) \\ & + 6a_3(1 - 2x + x^2 - 4it + 4ixt - 4t^2) + 6\bar{a}_3(-x^2 + 4ixt + 4t^2)\} \end{aligned}$$

and

$$\begin{aligned} \psi = & (9 - 36x + 72x^2 - 72x^3 + 72x^4 - 48x^5 + 16x^6) \\ & + 96t^2(3 + 3x - 4x^3 + 2x^4) + 384t^4(5 - 2x + 2x^2) + 1024t^6 \\ & + 24(a_3 + \bar{a}_3)(3x^2 - 2x^3 - 12t^2 + 24xt^2) \\ & + 48i(a_3 - \bar{a}_3)(3t + 6xt - 6x^2t + 8t^3) + 144a_3\bar{a}_3, \end{aligned}$$

and a_3 is a free complex parameter. We have found that the maximum of $|u(x, t, a_3)|$ is equal to 5, and it is obtained when

$$a_3 = -\frac{1}{12}.$$

At this a_3 value, the solution is

$$u_m(x, t) = 1 + \frac{\phi_m}{\psi_m}, \quad (4.5)$$

where

$$\begin{aligned} \phi_m = & 9 - 72\hat{x}^2 - 48\hat{x}^4 - 864t^2 - 3840t^4 - 1152\hat{x}^2t^2 \\ & + it(-180 - 288\hat{x}^2 + 192\hat{x}^4 + 384t^2 + 3072t^4 + 1536\hat{x}^2t^2), \end{aligned}$$




Figure 1. Second-order rogue waves with parameters: (a) $a_3 = -1/12$; (b) $a_3 = 5/3$; (c) $a_3 = -5i/2$; (d) $a_3 = 5i/2$. (Online version in colour.)

and

$$\begin{aligned} \psi_m = & \frac{9}{4} + 27\hat{x}^2 + 12\hat{x}^4 + 16\hat{x}^6 + 396t^2 + 1728t^4 + 1024t^6 - 288\hat{x}^2t^2 \\ & + 768\hat{x}^2t^4 + 192\hat{x}^4t^2, \end{aligned}$$

and $\hat{x} = x - 0.5$. This solution is displayed in figure 1*a*. It is easy to see that this solution is the special second-order rogue wave obtained by Akhmediev *et al.* (2009*a*) (after a shift in x). Thus, the special second-order rogue wave obtained by Akhmediev *et al.* is the one with the highest peak amplitude among all second-order rogue waves. At other a_3 values, however, we can obtain rogue waves which have very different solution dynamics from that in figure 1*a*. For instance, rogue waves at $a_3 = 5/3$, $-5i/2$ and $5i/2$ are displayed in figure 1*b–d*, respectively. In each of these solutions, three intensity humps appear at different times and/or space, and each intensity hump is roughly a first-order (Peregrine) rogue wave (4.2). Specifically, in figure 1*b*, the solution features double temporal bumps (elevations) at $x \approx -0.5$ and a single temporal bump at $x \approx 2.2$. In figure 1*c*, the solution first rises up and reaches a peak at $(x, t) \approx (0.5, -0.7)$. Afterwards, the solution temporally decays at $x \approx 0.5$, but two new bumps rise at the two sides. In figure 1*d*, the solution is similar to that in figure 1*c* but with a time reversal. Obviously, the rogue-wave dynamics in figure 1*b–d* are quite different from the one in figure 1*a*. The solution dynamics in figure 1*b–d* resemble those reported in Dubard *et al.* (2010), Dubard & Matveev (2011), Gaillard (2011) and Ankiewicz *et al.* (2011).




Figure 2. Third-order rogue waves with parameters (a_3, a_5) as: (a) $(-1/12, -1/240)$; (b) $(25/3, 0)$; (c) $(-25i/3, 0)$; (d) $(0, 50i/3)$. (Online version in colour.)

Next, we examine third-order rogue waves. In this case, we set $a_1 = a_2 = a_4 = 0$ without loss of generality. If one takes

$$a_3 = -\frac{1}{12} \quad \text{and} \quad a_5 = -\frac{1}{240},$$

then the corresponding solution $u_m(x, t)$ is equal to the third-order rogue wave obtained by Akhmediev *et al.* (2009a) except for a shift in x . This solution is displayed in figure 2a. The maximum amplitude of this solution is equal to 7, which occurs at $(x, t) = (1/2, 0)$. We have found that this special rogue-wave solution $u_m(x, t)$ is also the one with the highest peak amplitude among all third-order rogue waves $u(x, t; a_3, a_5)$. But if we take other (a_3, a_5) values, rogue waves with dynamics different from figure 2a will be obtained. Three such solutions, with $(a_3, a_5) = (25/3, 0)$, $(-25i/3, 0)$ and $(0, 50i/3)$, are displayed in figure 2b–d respectively. These solutions feature six intensity humps which appear at different times and/or space, and each intensity hump is roughly a first-order rogue wave (4.2). In figure 2b, the solution exhibits triple temporal bumps at $x \approx -2$, double temporal bumps at $x \approx 2$ and a single temporal bump at $x \approx 6$. In figure 2c, the solution develops a single hump first. Then this hump decays, but two new humps rise simultaneously at the two sides. Then these two humps decay, but

three additional humps develop simultaneously. In figure 2*d*, two intensity humps rise simultaneously at different spatial locations first. After they decay, additional four intensity humps arise at different locations and times. A remarkable feature in the rogue waves in figure 2*b–d* is the high regularity of their spatio-temporal patterns. For instance, the pattern in figure 2*c* is a highly symmetric triangle, while the one in figure 2*d* is like a pentagon. These spatio-temporal patterns of rogue waves are different from the ones reported in Dubard & Matveev (2011) and Gaillard (2011). The solution in figure 2*d* resembles the circular rogue waves obtained numerically in Kedziora *et al.* (2011), suggesting that those circular rogue waves are special cases of our general rogue-wave solutions.

The results shown above apparently can be extended to fourth- and higher order rogue waves. By special choices of the free parameters (a_3, a_5, a_7, \dots), we can reproduce the rogue waves obtained in Akhmediev *et al.* (2009*a*) as special cases. But other choices of those parameters can yield even richer spatio-temporal patterns, such as triangular patterns like figure 2*c*, but with more intensity humps such as 10, 15 and so on.

5. Summary and discussion

In this paper, we derived general N -th order rogue waves in the NLS equation by the bilinear method. These solutions were obtained from Gram determinant solutions of bilinear equations through dimension reduction and then further simplified to a very explicit form. We showed that these general rogue waves contain $N - 1$ free irreducible complex parameters. By different choices of these free parameters, we obtained rogue waves with novel spatio-temporal patterns. These new spatio-temporal patterns reveal the rich dynamics in rogue-wave solutions and deepen our understanding of the rogue-wave phenomena. We also showed that the rogue waves reported in Akhmediev *et al.* (2009*a*) are special solutions with the highest peak amplitude among all rogue waves of the same order.

We would like to point out that the new spatio-temporal patterns of rogue waves obtained in this paper may also find applications in other branches of applied mathematics and physics. For instance, the triangular rogue-wave patterns in figures 1*c* and 2*c* and their higher order extensions (with more intensity humps) closely resemble the spike pattern that forms after the point of gradient catastrophe in the semiclassical (zero-dispersion) limit of the NLS equation (see fig. 1 in Bertola & Tovbis 2010). The connection between these exact rogue-wave solutions and the semiclassical-NLS patterns is an interesting question which lies outside the scope of the present paper. For another instance, it has been shown recently that from rogue-wave solutions of the NLS equation, one can readily derive a subclass of smooth localized rational solutions to the Kadomtsev–Petviashvili (KP-I) equation (Dubard *et al.* 2010; Dubard & Matveev 2011). The application of our general higher order rogue waves in the NLS equation to the study of general higher order rational solutions in the KP-I equation is left for future studies.

The work of Y.O. is supported in part by JSPS Grant-in-Aid for Scientific Research (B-19340031, S-19104002) and for challenging Exploratory Research (22656026), and the work of J.Y. is supported

in part by the Air Force Office of Scientific Research (Grant USAF 9550-09-1-0228) and the National Science Foundation (Grant DMS-0908167).

Appendix A

In this appendix, we determine the number of free parameters in the rogue-wave solutions obtained in this paper. Since the solutions in theorem 2.1 are derived from the ones in lemma 3.2, we will examine solutions in lemma 3.2 below.

First we can factor out a_0 from A_i and b_0 from B_j . These factors cancel out in the formula (3.2), thus we will set $a_0 = b_0 = 1$ without loss of any generality.

Secondly, let us consider the effect of a constant shifting of (x_1, x_2) . By the shifting $(x_1, x_2) \rightarrow (x_1 + \alpha, x_2 + \beta)$, $m^{(n)}$ in (3.12) gets an exponential factor,

$$m^{(n)} \rightarrow m^{(n)} e^\theta \quad \text{and} \quad \theta = (p + q)\alpha + (p^2 - q^2)\beta, \quad (\text{A } 1)$$

consequently the (i, j) -component $A_i B_j m^{(n)}$ in (3.11) is also modified. Below we show that $A_i B_j m^{(n)}$ is modified as

$$A_i B_j m^{(n)} \rightarrow A_i B_j (m^{(n)} e^\theta) = e^\theta \hat{A}_i \hat{B}_j m^{(n)}, \quad (\text{A } 2)$$

where

$$\hat{A}_i = \sum_{k=0}^i \frac{\hat{a}_k}{(i-k)!} (p\partial_p)^{i-k}, \quad \hat{B}_j = \sum_{l=0}^j \frac{\hat{b}_l}{(j-l)!} (q\partial_q)^{j-l}, \quad (\text{A } 3)$$

and

$$\hat{a}_k = \sum_{\nu=0}^k a_\nu S_{k-\nu}(\mathbf{x}_0^+), \quad \mathbf{x}_0^+ = \left(p\alpha + 2p^2\beta, \frac{p\alpha + 4p^2\beta}{2}, \dots, \frac{p\alpha + 2^k p^2\beta}{k!}, \dots \right), \quad (\text{A } 4)$$

$$\hat{b}_l = \sum_{\nu=0}^l b_\nu S_{l-\nu}(\mathbf{x}_0^-), \quad \mathbf{x}_0^- = \left(q\alpha - 2q^2\beta, \frac{q\alpha - 4q^2\beta}{2}, \dots, \frac{q\alpha - 2^k q^2\beta}{k!}, \dots \right). \quad (\text{A } 5)$$

To prove (A 2), we notice that for the generator \mathcal{G} of the differential operators $(p\partial_p)^k$ defined by

$$\mathcal{G} = \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} (p\partial_p)^k = \exp(\lambda p\partial_p) = \exp(\lambda \partial_{\ln p}), \quad (\text{A } 6)$$

the relation

$$\mathcal{G}F(p, q) = F(e^\lambda p, q)$$

holds for any function $F(p, q)$. This relation is a special case of the previous relation (3.21). Thus,

$$e^{-\theta} \mathcal{G}(e^\theta F) = \exp((e^\lambda - 1)p\alpha + (e^{2\lambda} - 1)p^2\beta) \mathcal{G}F = \exp\left(\sum_{k=1}^{\infty} \frac{\lambda^k}{k!} (p\alpha + 2^k p^2\beta)\right) \mathcal{G}F,$$

whose coefficient of order λ^k gives

$$\frac{1}{k!} (p\partial_p)^k (e^\theta F) = e^\theta \sum_{\nu=0}^k S_\nu(\mathbf{x}_0^+) \frac{1}{(k-\nu)!} (p\partial_p)^{k-\nu} F.$$

Similarly, we have

$$\frac{1}{l!} (q\partial_q)^l (e^\theta F) = e^\theta \sum_{\nu=0}^l S_\nu(\mathbf{x}_0^-) \frac{1}{(l-\nu)!} (q\partial_q)^{l-\nu} F.$$

Therefore,

$$\begin{aligned} A_i B_j (m^{(n)} e^\theta) &= \sum_{k=0}^i \sum_{l=0}^j a_k b_l \frac{1}{(i-k)!} (p\partial_p)^{i-k} \frac{1}{(j-l)!} (q\partial_q)^{j-l} (m^{(n)} e^\theta) \\ &= e^\theta \sum_{k=0}^i \sum_{l=0}^j a_k b_l \sum_{\mu=0}^{i-k} S_\mu(\mathbf{x}_0^+) \frac{1}{(i-k-\mu)!} (p\partial_p)^{i-k-\mu} \\ &\quad \times \sum_{\nu=0}^{j-l} S_\nu(\mathbf{x}_0^-) \frac{1}{(j-l-\nu)!} (q\partial_q)^{j-l-\nu} m^{(n)} \\ &= e^\theta \sum_{k=0}^i \sum_{l=0}^j \frac{\hat{a}_k}{(i-k)!} (p\partial_p)^{i-k} \frac{\hat{b}_l}{(j-l)!} (q\partial_q)^{j-l} m^{(n)} = e^\theta \hat{A}_i \hat{B}_j m^{(n)}, \end{aligned}$$

which proves equation (A 2).

Now we take $p = q = 1$. Then from equations (A 4)–(A 5), we get

$$\hat{a}_0 = a_0 = 1, \quad \hat{a}_1 = a_1 + \alpha + 2\beta, \dots, \quad \hat{b}_0 = b_0 = 1, \quad \hat{b}_1 = b_1 + \alpha - 2\beta, \dots$$

Thus by a shifting of $(x_1, x_2) \rightarrow (x_1 + \alpha, x_2 + \beta)$ with $\alpha = -(a_1 + b_1)/2$ and $\beta = -(a_1 - b_1)/4$, we obtain $\hat{a}_1 = \hat{b}_1 = 0$. When this shifting is combined with shifts of higher coefficients $a_2 \rightarrow \hat{a}_2, a_3 \rightarrow \hat{a}_3, \dots, b_2 \rightarrow \hat{b}_2, b_3 \rightarrow \hat{b}_3, \dots$, the solution τ_n depends on parameters $(\hat{a}_2, \hat{a}_3, \dots; \hat{b}_2, \hat{b}_3, \dots)$ only. In other words, by a shift of (x_1, x_2) , we can normalize $a_1 = b_1 = 0$.

Thirdly, from the expressions of m_{ij}^n in (3.11) and the expressions of A_i and B_j , we see that in the determinant formula for τ_n in (3.13), when we subtract the product of the first row and a_2 from the second row, and subtract the product of the second row and a_2 from the third row, ..., and subtract the product of the i th row and a_2 from the $(i+1)$ -th row, and then subtract the product of the first column and b_2 from the second column, and subtract the product of the second column and b_2 from the third column, etc., we can remove the parameter a_2 and b_2 from the solution formula (3.13). By similar treatments, we can remove all other even coefficients a_4, a_6, \dots and b_4, b_6, \dots as well. In other words, we can set $a_2 = a_4 = a_6 = \dots = 0$ and $b_2 = b_4 = b_6 = \dots = 0$ without any loss of generality.

By summarizing the above results, we see that without any loss of generality, we can set

$$a_0 = b_0 = 1, \quad a_2 = a_4 = a_6 = \dots = b_2 = b_4 = b_6 = \dots = 0.$$

In addition, by a shift of (x_1, x_2) , we can normalize $a_1 = b_1 = 0$. Combined with the complex conjugacy condition $b_k = \bar{a}_k$ in (3.17), we then find that the N -th order rogue-wave solutions in theorem 2.1 have $N - 1$ free irreducible complex parameters, $a_3, a_5, \dots, a_{2N-1}$.

References

- Ablowitz, M. J. & Herbst, B. M. 1990 On homoclinic structure and numerically induced chaos for the nonlinear Schrödinger equation. *SIAM J. Appl. Math.* **50**, 339–351. (doi:10.1137/0150021)
- Akhmediev, N., Eleonskii, V. M. & Kulagin, N. E. 1985 Generation of a periodic sequence of picosecond pulses in an optical fiber: exact solutions. *Sov. Phys. JETP* **89**, 1542–1551. [In Russian.]
- Akhmediev, N., Eleonskii, V. M. & Kulagin, N. E. 1988 Exact first-order solutions of the nonlinear Schrödinger equation. *Theor. Math. Phys.* **72**, 809–818. (doi:10.1007/BF01017105)
- Akhmediev, N., Ankiewicz, A. & Soto-Crespo, J. M. 2009a Rogue waves and rational solutions of the nonlinear Schrödinger equation. *Phys. Rev. E* **80**, 026 601. (doi:10.1103/PhysRevE.80.026601)
- Akhmediev, N., Ankiewicz, A. & Taki, M. 2009b Waves that appear from nowhere and disappear without a trace. *Phys. Lett. A* **373**, 675–678. (doi:10.1016/j.physleta.2008.12.036)
- Akhmediev, N., Soto-Crespo, J. M. & Ankiewicz, A. 2009c Extreme waves that appear from nowhere: on the nature of rogue waves. *Phys. Lett. A* **373**, 2137–2145. (doi:10.1016/j.physleta.2009.04.023)
- Ankiewicz, A., Kedziora, D. J. & Akhmediev, N. 2011 Rogue wave triplets. *Phys. Lett. A* **375**, 2782–2785. (doi:10.1016/j.physleta.2011.05.047)
- Benney, D. J. & Newell, A. C. 1967 Nonlinear wave envelopes. *J. Math. Phys.* **46**, 133–139.
- Bertola, M. & Tovbis, A. 2010 Universality for the focusing nonlinear Schrödinger equation at the gradient catastrophe point: rational breathers and poles of the tritronquee solution to Painlevé I. See <http://arxiv.org/abs/1004.1828>.
- Dubard, P. & Matveev, V. B. 2011 Multi-rogue waves solutions to the focusing NLS equation and the KP-I equation. *Nat. Hazards Earth. Syst. Sci.* **11**, 667–672. (doi:10.5194/nhess-11-667-2011)
- Dubard, P., Gaillard, P., Klein, C. & Matveev, V. B. 2010 On multi-rogue wave solutions of the NLS equation and positon solutions of the KdV equation. *Eur. Phys. J. Spec. Top.* **185**, 247–258. (doi:10.1140/epjst/e2010-01252-9)
- Gaillard, P. 2011 Families of quasi-rational solutions of the NLS equation and multi-rogue waves. *J. Phys. A Math. Theor.* **44**, 435204. (doi:10.1088/1751-8113/44/43/435204)
- Hasegawa, A. & Tappert, F. 1973 Transmission of stationary nonlinear optical pulses in dispersive dielectric fibers. *Appl. Phys. Lett.* **23**, 142. (doi:10.1063/1.1654836)
- Hirota, R. 2004 *The direct method in soliton theory*. Cambridge, UK: Cambridge University Press. (doi:10.1017/CBO9780511543043)
- Ikeda, T. & Yamada, H.-F. 2002 Polynomial τ -functions of the NLS-Toda hierarchy and the Virasoro singular vectors. *Lett. Math. Phys.* **60**, 147–156. (doi:10.1023/A:1016167008456)
- Its, A. R., Rybin, A. V. & Salle, M. A. 1988 Exact integration of nonlinear Schrödinger equation. *Theor. Math. Phys.* **74**, 29–45. (doi:10.1007/BF01018207)
- Jimbo, M. & Miwa, T. 1983 Solitons and infinite dimensional Lie algebras. *Publ. RIMS Kyoto Univ.* **19**, 943–1001. (doi:10.2977/prims/1195182017)
- Kedziora, D. J., Ankiewicz, A. & Akhmediev, N. 2011 Circular rogue wave clusters. *Phys. Rev. E* **84**, 056611. (doi:10.1103/PhysRevE.84.056611)

- Kibler, B., Fatome, J., Finot, C., Millot, G., Dias, F., Genty, G., Akhmediev, N. & Dudley, J. M. 2010 The Peregrine soliton in nonlinear fibre optics. *Nat. Phys.* **6**, 790–795. (doi:10.1038/nphys1740)
- Ohta, Y., Wang, D. & Yang, J. 2011 General N-dark-dark solitons in the coupled nonlinear Schrödinger equations. *Stud. Appl. Math.* **127**, 345–371. (doi:10.1111/j.1467-9590.2011.00525.x)
- Peregrine, D. H. 1983 Water waves, nonlinear Schrödinger equations and their solutions. *J. Aust. Math. Soc. B* **25**, 16–43. (doi:10.1017/S0334270000003891)
- Sato, M. 1981 Soliton equations as dynamical systems on a infinite dimensional Grassmann manifolds. *RIMS Kokyuroku* **439**, 30–46.
- Solli, D. R., Ropers, C., Koonath, P. & Jalali, B. 2007 Optical rogue waves. *Nature* **450**, 1054–1057. (doi:10.1038/nature06402)
- Zakharov, V. E. 1968 Stability of periodic waves of finite amplitude on the surface of a deep fluid. *J. Appl. Mech. Tech. Phys.* **9**, 190–194. (doi:10.1007/BF00913182)
- Zakharov, V. E. & Shabat, A. B. 1972 Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in nonlinear media. *Sov. Phys. JETP* **34**, 62–69.